Effective Treatment of Invasive Aspergillus fumigatus Infection Using Combinations of Topical and Systemic Antifungals in a Severely Burned Patient

Anne Fournier^{1,8}, Olivier Pantet², Samia Guerid³, Philippe Eggimann², Jean-Luc Pagani², Jean-Pierre Revelly², Philippe M. Hauser⁴, Oscar Marchetti⁵, Sara Fontanella⁶, Igor Letovanec⁶, François Ravat⁷, Mette M. Berger², André Pannatier^{1,8}, Pierre Voirol^{1,8}, Yok-Ai Que²

¹Service of Pharmacy, ²Service of Intensive Care Medicine, ³Service of Plastic and Reconstructive Surgery, ⁴Institute of Microbiology, ⁵Service of Infectious Diseases, and ⁶Service of Pathology, Centre Hospitalier Universitaire Vaudois and University of Lausanne, Switzerland; ⁷Centre des Brûlés, Centre Hospitalier St. Joseph et St Luc, Lyon, France; and ⁸School of PharmaceuticalSciences, University of Geneva, University of Lausanne.

Abstract

We describe an invasive Aspergillus fumigatus deep-burn wound infection in a severely burned patient that was successfully treated with a combination of topical terbinafine and systemic voriconazole antifungal therapy. To our knowledge, this is the first case report describing the effective control of an invasive deep-burn wound infection using this combination.

Published in : J Burn Care Res 2014;

doi: 10.1097/BCR.00000000000143

Contact: <u>anne.henry@chuv.ch</u>